arXiv:1603.08639v1 [math.DS] 29 Mar 2016

Fast growth of the number of periodic points in generic families of two-dimensional real-analytic area-preserving diffeomorphisms

Masayuki Asaoka

March 30, 2016

Abstract

We prove that there exists an open subset of the set of real-analytic area-preserving diffeomorphisms of a closed surface in which generic fimite-parapeter families exhibit fast growth of the number of periodic points for any parameter.

1 Introduction

Growth rate of the number of periodic points has been interested in people working on dynamical systems for long time. If the number of periodic points grows at most exponentially (we say that the map exhibits at most exponential growth in this case), then the dynamical zeta function of the map has positive convergent radius.

In 1965, Artin and Mazur [1] proved that C^r maps which exhibits at most exponential growth are dense in the space of C^r-maps on a closed manifold. For one-dimensional dynamics, any C^∞ map on a compact interval without ∞-flat critical points exhibits at most exponential growth. This is an easy corollary of a result of Martens, de Melo, and van Strien [12] (see also [8, Corollary 1]). In particular, any C^ω map on a compact interval exhibits at most exponential growth. If we drop the flatness of critical point, superexponential growth is possible. In fact, for any given sequence of positive integers and any $1 \leq r < \infty$, Kaloshin and Kozlovski [8] constructed an example of C^r unimodal map on the unit interval whose number of periodic points grows faster than the sequence[1] (in this case, we say that the map exhibits fast growth of the number of periodic points). For higher dimensional case, Kaloshin [7] showed that fast growth is generic in the Newhouse domain in the space of C^r surface diffeomorphisms for any $2 \leq r \leq \infty$[2]. Kaloshin and Saprykina [9] and the author, Shinohara, and Turaev [2] also gave open sets of maps in which generic maps exhibit fast

[1] In the appendix of this paper, we give a simple example for C^∞ case.

[2] Kaloshin stated his result for finite r. However, his proof works for C^∞ case.

growth in other settings. However, as far as the author's knowledge, there was no known C^ω example which exhibits fast growth of the number of periodic points.[3] In this paper, we show that fast growth is generic in an open subset of the set of C^r area-preserving diffeomorphisms on a closed surface for $r = \infty$ or ω.

Moreover, we will show that fast growth of the number of periodic points is typical in the sense of Kolmogorov-Arnold in this open subset. Let Δ_k be the k-dimensional unit closed ball centered at the origin. A property of a C^r map ($1 \leq r \leq \infty$ or $r = \omega$) is *C^r-typical* in an open subset $\mathcal{U}$ of C^r area-preserving maps on a manifold if for any $k \geq 1$ and any generic C^r family $(f_\delta)_{\delta \in \Delta_k}$ of maps in $\mathcal{U}$, the property holds for Lebesgue almost all $\delta \in \Delta_k$. See [6] for more information on typicalness and related concepts. In many cases, typicalness is harder to show than genericity. For example, finiteness of periodic attractor and at most exponential growth of the numper of periodic points for C^r-typical diffeomorphisms have been a long-standing open problems. Recently, Berger [3] proved that finiteness of periodic attractors is not typical for three-dimensional diffeomorphisms. At most exponential growth for typical diffeomorphism is still open. However, the main theorem of this paper implies that fast growth is C^r-typical in an open subset of the set of C^r area-preserving diffeomorphisms on a closed surface for $r = \infty$ or ω.

In the rest of the article, we fix $r = \infty$, or ω. For C^r compact manifolds M_1 and M_2, let $C^r(M_1, M_2)$ be the space of C^r maps from M_1 to M_2 with the C^r topology. See [11] for the definition of C^ω topology. It is known that $C^r(M_1, M_2)$ are metrizable as a complete metric space. In particular, it is a Baire space.

Let Σ be a C^r closed orientable two-dimensional manifold and μ be a C^r volume form of Σ. By $\mathrm{Diff}^r_\mu(\Sigma)$, we denote the group of C^r-diffeomorphisms of Σ which preserve the volume form μ. This group admits the C^r-topology as a subset of $C^r(\Sigma, \Sigma)$. Recall that Δ_k is the k-dimensional closed unit disk centered at th origin. We say that a family $(h_\delta)_{\delta \in \Delta_k}$ of maps in $C^r(M_1, M_2)$ is of class C^r if the map $(x, \delta) \mapsto h_\delta(x)$ from $M_1 \times \Delta_k$ to M_2 is of class C^r. The set of C^r families $(f_\delta)_{\delta \in \Delta_k}$ of maps in $C^r(M_1, M_2)$ is endowed with the C^r topology as a subset of $C^r(M_1 \times \Delta_k, M_2)$.

For an elliptic fixed point p of a diffeomorphism $f \in \mathrm{Diff}^r_\mu(\Sigma)$, the Birkhoff normal form of f at p up to the third order is

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + O((x^2+y^2)^2),$$

where $\alpha(x, y) = \alpha_0 + \alpha_1(x^2+y^2)$ with real numbers α_0, α_1 (see *e.g.* [13, Theorem 2.11]). We say that the elliptic fixed point p satisfies *the Moser stability condition* if $\alpha_1 > 0$. For $f \in \mathrm{Diff}^r_\mu(\Sigma)$, let $\mathrm{Fix}_h(f)$ (*resp.* $\mathrm{Fix}_e(f)$) be the set of hyperbolic (*resp.* elliptic) fixed points. By $\#S$, we denote the cardinality of a set S. The following is the main theorem of this paper.

[3] A related example is given by Kozlovski[10]

Theorem 1.1. *Suppose that a diffeomorphism $f_0 \in \mathrm{Diff}^r_\mu(\Sigma)$ has an elliptic fixed point with the Moser stability condition. Then, there exists an open neighborhood $\mathcal{U}$ of f_0 such that for any sequence $(k_n)_{n\geq 1}$ of positive integers, a generic family $(f_\delta)_{\delta\in\Delta_k}$ of diffeomorphisms in $\mathcal{U}$ satisfies that*

$$\limsup_{n\to\infty} \frac{\min\{\#\,\mathrm{Fix}_h(f_\delta^n), \#\,\mathrm{Fix}_e(f_\delta^n)\}}{a_n} = \infty \tag{1}$$

for any $\delta \in \Delta_k$. In particular, the condition (1) is C^r-typical in $\mathcal{U}$.

2 Proof of Theorem 1.1

We fix $k \geq 1$ and let Δ be the closed unit ball of dimension k. By $\mathrm{Diff}^r_\mu(\Sigma;\Delta)$ be the set of C^r family $(f_\delta)_{\delta\in\Delta}$ of diffeomorphisms in $\mathrm{Diff}^r_\mu(\Sigma)$. We say that a family $(f_\delta)_{\delta\in\Delta}$ in $\mathrm{Diff}^r_\mu(\Sigma;\Delta)$ admits a KAM circle if there exists a family $(\psi_\delta)_{\delta\in\Delta}$ of C^r embeddings from $S^1\times(-1,1)$ to Σ and a real number α such that

$$f_\delta \circ \psi_\delta(\theta, 0) = \psi_\delta(\theta+\alpha, 0)$$

and

$$\left(\frac{\partial}{\partial r}\left[\pi_\theta \circ (\psi_\delta^{-1}\circ f_\delta \circ \psi_\delta)\right]\right)(\theta, 0) > 0$$

for any $\delta\in\Delta$ and $\theta \in S^1$, where $\pi_r : S^1\times(-1,1)\to(-1,1)$ is the projection to the second coordinate. The former condition means that f_δ is C^r conjugate to the rigid rotation of angle α on an invariant circle $\psi_\delta(S^1\times\{0\})$. The latter means that f_δ 'twists' on a neighborhood of the circle. The following is an easy consequence of a classical result by Moser (see [14, §32-34] for $r=\omega$ and [13, Theorem 2.11] for $r=\infty$).

Proposition 2.1. *Let f_0 be a diffeomorphism in $\mathrm{Diff}^r_\mu(\Sigma)$ and p be an elliptic fixed point of f_0 with the Moser stability condition. Then, there exists a neighborhood $\mathcal{U}$ of f_0 in $\mathrm{Diff}^r_\mu(\Sigma)$ such that any C^r family $(f_\delta)_{\delta\in\Delta}$ of elements in $\mathcal{U}$ admits a KAM circle.*

The following proposition implies Theorem 1.1.

Proposition 2.2. *Suppose that a family $(f_\delta)_{\delta\in\Delta}$ in $\mathrm{Diff}^r_\mu(\Sigma;\Delta)$ admits a KAM circle. Then for any neighborhood $\mathcal{V}$ of $(f_\delta)_{\delta\in\Delta}$ in $\mathrm{Diff}^r_\mu(\Sigma;\Delta)$, any sequence $(a_n)_{n\geq1}$ of positive integers, and any $n_0\geq 1$ there exists a family $(\tilde f_\delta)_{\delta\in\Delta}$ in $\mathcal{V}$ and an integer $m\geq n_0$ such that*

$$\min\{\#\,\mathrm{Fix}_h(\tilde f_\delta^m), \#\,\mathrm{Fix}_e(\tilde f_\delta^m)\} \geq a_m$$

for any $\delta\in\Delta$.

Proof of Theorem 1.1 from Proposition 2.2. Let $\mathcal{U}$ be an open neighborhood of f_0 in $\mathrm{Diff}^r_\mu(\Sigma)$ given by Proposition 2.1. Fix a sequence $(a_n)_{n\geq1}$ of positive integers. Let $\mathcal{U}_\Delta$ be the set of C^r families $(f_\delta)_{\delta\in\Delta}$ with $f_\delta\in\mathcal{U}$ for any δ. Put

$$\mathcal{V}_m = \{(f_\delta)_{\delta\in\Delta}\in\mathcal{U}_\Delta \mid \min\{\#\,\mathrm{Fix}_h(f_\delta^m), \#\,\mathrm{Fix}_e(f_\delta^m)\}\geq a_m \text{ for any } \delta\in\Delta\}.$$

By the persistence of hyperbolic and elliptic fixed points, the set $\mathcal{V}_m$ is open for any $m \geq 1$. By Proposition 2.2, the set $\bigcup_{m \geq n} \mathcal{V}_m$ is dense in $\mathcal{U}_\Delta$ for any $n \geq 1$. The set $\mathcal{R} = \bigcap_{n \geq 0} \bigcup_{m \geq n} \mathcal{V}_m$ is residual in $\mathcal{U}_\Delta$ and any family $(f_\delta)_{\delta \in \Delta}$ in $\mathcal{R}$ satisfies (1). □

The rest of this section is devoted to the proof of Proposition 2.2.

Lemma 2.3. *Let M be a C^r compact submanifold of $\mathbb{R}^m \times \Delta$. For any C^r function u on a neighborhood of M and any $s \geq 1$, there exists a C^r function $\bar{u}$ on $\mathbb{R}^n \times \Delta$ such that $j^s_x \bar{u} = j^s_x u$ for any $x \in M$, where $j^s_x u$ is the s-jet of u at x.*

Proof. For the case $r = \infty$, the lemma follows from the standard argument using a bump function.

Let us prove the lemma for the case $r = \omega$. Take a C^ω function u on a neighborhood of M. We can find a k-dimensional closed ball Δ' of radius greater than one and centered at the origin such that M extends to a compact submanifold M' of $\mathbb{R}^m \times \Delta'$ and u extends to a C^ω function u_h on a neighborhood V_u of M'. Let $\mathcal{A}$ be the sheaf of C^ω functions on $\mathbb{R}^m \times \operatorname{Int} \Delta'$ and $\mathcal{F}$ be the subsheaf of $\mathcal{A}$ defined by

$$\mathcal{F}(U) = \{u \in \mathcal{A}(U) \mid j^s_x u = 0 \text{ for any } x \in M\}.$$

It is easy to see that the sheaf $\mathcal{F}$ is of finite type. This means the sheaf $\mathcal{F}$ is coherent since it is a subsheaf of a coherent sheaf $\mathcal{A}$. Since $\mathbb{R}^m \times \operatorname{Int} \Delta'$ is C^ω diffeomorphic to $\mathbb{R}^{m+k}$, we have $H^1(\mathbb{R}^m \times \operatorname{Int} \Delta', \mathcal{F}) = \{0\}$ by Cartan's Theorem B. A short exact sequence $0 \to \mathcal{F} \to \mathcal{A} \to \mathcal{A}/\mathcal{F} \to 0$ of sheaves induces a exact sequence

$$H^0(\mathbb{R}^m \times \operatorname{Int} \Delta', \mathcal{A}) \to H^0(\mathbb{R}^m \times \operatorname{Int} \Delta', \mathcal{A}/\mathcal{F}) \to H^1(\mathbb{R}^m \times \operatorname{Int} \Delta', \mathcal{F}) = \{0\}.$$

of the cohomology of sheaves. The C^ω extension u_h of u to V_u defines an element $[u_h]$ of $H^0(\mathcal{A}/\mathcal{F})$ by $[u_h]_x = (u_h)_x$ if $x \in M'$ and $[u_h]_x = 0$ if $x \notin M'$, where $(u_h)_x$ means the germ in the stalk $\mathcal{A}_x$ represented by u_h at x. By the above exact sequence, there exists a C^ω function $\bar{u}$ on $\mathbb{R}^m \times \operatorname{Int} \Delta'$ such that $j^s_x(\bar{u} - u_h) = 0$ for any $x \in M'$. □

Lemma 2.4. *There exists a C^r family $(H_\delta)_{\delta \in \delta}$ of functions on Σ such that the Hamiltonian vector field X_δ generated by H_δ with respect to the 2-form μ satisfies that*

$$D\psi_\delta^{-1}(X_\delta)_{(\theta,0)} = \frac{\partial}{\partial \theta}. \tag{2}$$

Proof. The set

$$C_\Delta = \{(\psi_\delta(\theta, 0), \delta) \mid \theta \in S^1, \delta \in \Delta\}$$

is a C^r compact submanifold $\Sigma \times \Delta$. We embed Σ into $\mathbb{R}^m$ with large m as a C^r submanifold and identify C_Δ with a C^r compact submanifold of $\mathbb{R}^m \times \Delta$. Define a C^r family $(\beta_\delta)_{\delta \in \Delta}$ of functions on $S^1 \times (-1, 1)$ by $\psi_\delta^* \mu = \beta_\delta \cdot d\theta \wedge dr$. We can find a C^r function u on an open neighborhood of C_Δ in $\mathbb{R}^m \times \Delta$ such

that $u(\psi_\delta(\theta,r),\delta) = r\cdot\beta_\delta(\theta,r)$ for any (θ,r) in a small neighborhood of $S^1\times\{0\}$ and any $\delta\in\Delta$. By Lemma 2.3, there exists a C^r function H on $\mathbb{R}^m\times\Delta$ such that $dH_x = du_x$ for any $x\in C_\Delta$. Define a C^r family of functions on Σ by $H_\delta(x) = H(x,\delta)$ for $(x,\delta)\in\Sigma\times\Delta\subset\mathbb{R}^m\times\Delta$. Then,

$$d(H_\delta\circ\psi_\delta)_{(\theta,0)} = d(r\cdot\beta_\delta)_{(\theta,0)} = (\beta_\delta dr)_{(\theta,0)}$$

for any $\theta\in S^1$. This implies that

$$d(H\circ\psi_\delta)(X) = \beta_\delta(\theta,0)dr(X) = \psi_\delta^*\mu\left(\frac{\partial}{\partial\theta}, X\right)$$

for any $X\in T_{(\theta,0)}(S^1\times(-1,1))$. Therefore, the Hamiltonian vector field X_δ of H_δ satisfies (2). □

Lemma 2.5. *For any integers $k,l\geq 1$, there exists a C^r family $(G_\delta)_{\delta\in\Delta}$ of functions on Σ such that Hamiltonian flow $(\Psi_\delta^t)_{t\in\mathbb{R}}$ associated with G_δ with respect to the 2-form μ satisfies $\Psi_\delta^t\circ\psi_\delta(j/2kl,0) = \psi_\delta(j/2kl)$ and*

$$D(\psi_\delta^{-1}\circ\Psi_\delta^t\circ\psi_\delta)_{(j/2kl,0)} = \begin{cases} \begin{pmatrix}\cos t & -\sin t\\ \sin t & \cos t\end{pmatrix} & (j=0,\dots,l-1)\\ \begin{pmatrix}\cos t & \sin t\\ -\sin t & \cos t\end{pmatrix} & (j=l,\dots,2l-1)\\ \begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix} & (j=2l,\dots,2kl-1)\end{cases}$$

for any $\delta\in\Delta$ and $j=0,\dots,2kl-1$.

Proof. Similar to Lemma 2.4, we apply Lemma 2.3 to the one-dimensional submanifold

$$\{(\psi_\delta(j/2kl,0),\delta)\mid j=0,\dots,2kl-1,\delta\in\Delta\}$$

of $\Sigma\times\Delta$. Then, we obtain a C^r family G_δ of functions on Σ such that

$$j^2_{(j/2kl,0)}(G_\delta\circ\psi_\delta)(\theta,r) = \begin{cases}(\theta-(j/2kl))^2+r^2 & (j=0,\dots,l-1)\\ -(\theta-(j/2kl))^2-r^2 & (j=l,\dots,2l-1)\\ 0 & (j=2l,\dots,2kl-1).\end{cases}$$

The point $\psi_\delta(j/2kl,0)$ is a critical point of G_δ, and hence, it is a stationary point of the Hamiltonian flow Ψ_δ^t generated by G_δ. At the stationary point of a Hamiltonian flow, the derivative of the flow is determined by the 2-jet of the Hamiltonian. By a direct computation, we can check the required condition on the derivative of $\psi_\delta^{-1}\circ\Psi_\delta^t\circ\psi_\delta$ at $(j/2kl,0)$. □

Now, we prove Proposition 2.2. Fix an open neighborhood $\mathcal{V}$ of $(f_\delta)_{\delta\in\Delta}$, a sequence $(a_n)_{n\geq 1}$ of positive integers, and $n_0\geq 1$. By Lemma 2.4, there exists a C^r family $(H_\delta)_{\delta\in\Delta}$ of functions on Σ such that the Hamiltonian vector field

X_δ satisfies $D\psi_\delta^{-1}(X_\delta)_{(\theta,0)} = \frac{\partial}{\partial\theta}$ for any $\theta \in S^1$. Let $\Phi_\delta = (\Phi_\delta^t)$ be the flow generated by X_δ. It preserves μ and satisfies that

$$(\Phi_\delta^t \circ f_\delta)(\psi_\delta(\theta, 0)) = \psi_\delta(\theta + \alpha + t, 0).$$

Take small $T > 0$ so that $\alpha + T$ is a rational number with denominator $\geq n_0$, the family $(\Phi_\delta^T \circ f_\delta)_{\delta\in\Delta}$ is contained in $\mathcal{V}$, and this family satisfies

$$\tau_\delta(\theta) := \frac{\partial}{\partial r}[\pi_\theta \circ \psi_\delta^{-1} \circ (\Phi_\delta^T \circ f_\delta) \circ \psi_\delta](\theta, 0) > 0$$

for any $\delta \in \Delta$ and $\theta \in S^1$. Let k be the denominator of $\alpha + T$. Then, any point of $\psi_\delta(S^1 \times \{0\})$ is a fixed point of $(\Phi_\delta^T \circ f_\delta)^k$ and

$$D(\psi_\delta^{-1} \circ \Phi_\delta^T \circ f_\delta \circ \psi_\delta)_{(\theta,0)} = \begin{pmatrix} 1 & \tau_\delta(\theta) \\ 0 & * \end{pmatrix}.$$

Since f_δ preserves the 2-form μ, we have

$$D(\psi_\delta^{-1} \circ (\Phi_\delta^T \circ f_\delta)^k \circ \psi_\delta)_{(\theta,0)} = \begin{pmatrix} 1 & \tau_\delta^k(\theta) \\ 0 & 1 \end{pmatrix},$$

where $\tau_\delta^k(\theta)$ is a linear combination of $\tau_\delta(\theta), \tau_\delta(\theta + 1/k), \ldots, \tau_\delta(\theta + (k-1)/k)$ with positive coefficients. In particular, $\tau_\delta^k(\theta)$ is positive.

By Lemma 2.5 with $(k, l) = (k, a_k)$, there exists a C^r family $(G_\delta)_{\delta\in\Delta}$ of functions on Σ such that the Hamiltonian flow $(\Psi_\delta^t)_{\delta\in\Delta}$ generated by G_δ has the derivative given in the lemma. Then, $\psi_\delta(j/2ka_k, 0)$ is a fixed point of $(\Psi_\delta^t \circ \Phi_\delta^T \circ f_\delta)^k$ and

$$\begin{aligned} &D(\psi_\delta^{-1} \circ (\Psi_\delta^t \circ \Phi_\delta^T \circ f_\delta)^k \circ \psi_\delta)_{(j/2ka_k,0)} \\ &= \begin{cases} \begin{pmatrix} 1 & \tau_\delta^k(\theta) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} & (j = 0, \ldots, a_k - 1) \\ \begin{pmatrix} 1 & \tau_\delta^k(\theta) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} & (j = a_k, \ldots, 2a_k - 1). \end{cases} \end{aligned}$$

for any $j = 0, \ldots, 2a_k$ and $\delta \in K$. Since $\tau_\delta(\theta)$ is a positive function, the fixed point $\psi_\delta(j/2ka_k, 0)$ is hyperbolic if $j = 0, \ldots, a_k - 1$ and elliptic if $j = a_k, \ldots, 2a_k - 1$ for any sufficiently small $t > 0$ and $\delta \in \Delta$. The family $(\Psi_\delta^t \circ \Phi_\delta^T \circ f_\delta)_{\delta\in\Delta}$ is contained in $\mathcal{V}$ if $t > 0$ is small, and this family satisfies the the required conditions.

A Another example of smooth one-dimensional dynamics with fast growth

In this appendix, we give a simple example of a C^∞ map on the unit interval which exhibits fast growth of the number of periodic points.

Theorem A.1. *For any sequence $(a_n)_{n\geq 1}$ of positive integers, there exists a C^∞ self-map f on the unit interval such that*

$$\limsup_{n\to\infty} \frac{\#\operatorname{Fix}(f^n)}{a_n} = \infty. \tag{3}$$

As mentioned in Section 1, such a map must have a flat critical point by Martens, de Melo, and van Strien's theorem [12]. For any finite $r \geq 1$, Kaloshin and Kozlovski[8] constructed a C^r unimodal map with the above property which is of class C^∞ except at the critical point.

Proof. Our example is based on de Melo's construction of C^∞ interval map with infinitely many periodic attractors [5].

Recall that the C^∞ topology of the space of C^∞ self-maps on $[0,1]$ is given by the metric

$$d_{C^\infty}(f,g) = \sum_{s=0}^{\infty} 2^s \frac{\|f^{(s)} - g^{(s)}\|_\infty}{1 + \|f^{(s)} - g^{(s)}\|_\infty},$$

where $\|h\|_\infty = \sup_{x\in[0,1]} |h(x)|$.

Fix $0 < \delta < 1/6$. Let $f_0 : [0,1] \to [0,1]$ be a C^∞ map which satisfies that

$$f_0(x) = \begin{cases} 2x & \left(0 \leq x \leq \frac{1}{3}\right) \\ 1 & \left(\frac{1}{2} - \delta \leq x \leq \frac{1}{2} + \delta\right) \\ 2(1-x) & \left(\frac{2}{3} \leq x \leq 1\right). \end{cases}$$

We put

$$\begin{aligned} y_k &= \frac{1}{2} + 6^{-k} \cdot 4\delta, \\ J_k &= [y_k - 6^{-k}\delta, y_k + 6^{-k}\delta], \\ I_k &= [y_k - 6^{-k} \cdot 2\delta, y_k + 6^{-k} \cdot 2\delta]. \end{aligned}$$

Notice that I_k and $I_{k'}$ are disjoint if $k \neq k'$. For $k \geq 1$, we take a C^∞ function g_k on $[0,1]$ such that $0 \leq g_k \leq 1$ and

$$g_k(x) = \begin{cases} 0 & (x \in [0,1] \setminus I_k) \\ x & (x \in J_k). \end{cases}$$

Fix $\epsilon > 0$. We construct a C^∞ map $f : [0,1] \to [0,1]$ with the condition (3) such that $d_{C^\infty}(f, f_0) < \epsilon$. For $k \geq 1$, we can choose $n_k \geq 1$ such that $d_{C^\infty}(2^{-n_k} g_k, 0) < 2^{-k+1}\epsilon$. Define a map $f_1 : [0,1] \to [0,1]$ by

$$f = f_0 - \sum_{k=1}^{\infty} 2^{-n_k} g_k.$$

The map f_1 is of class C^∞ and it satisfies $d_{C^\infty}(f_1, f_0) \leq \epsilon/2$. For $x \in J_k$, we have

$$f^{n_k}(x) = f^{n_k - 1}(2^{-n_k+1} g_k(x)) = g_k(x) = x$$

For each $k \geq 1$, there exists a function h_k on $[0,1]$ such that $h(x) = 0$ if $x \in [0,1] \setminus J_k$, the set $\{x \in J_k \mid h(x) = 0\}$ contains $k \cdot a_{n_k}$ points, and $d_{C^\infty}(h_k) < 2^{-k}\epsilon$. Then, the map

$$f = f_0 - \sum_{k=1}^{\infty} (2^{-n_k} g_k + h_k)$$

is of class C^∞ which satisfies the condition (3) and $d_{C^\infty}(f, f_0) < \epsilon$. □

To simplify the construction, we started with a specific map f_0. However, the above perturbation method can be generalized to any one-dimensional map with a flat critical point which is a homoclinic point of a repelling periodic point since any flat critical point can be perturbed into a locally constant critical point and any one-dimensional map can be linearized at repelling periodic point.

References

[1] M. Artin and B. Mazur, On periodic points, Ann. of Math., **81** (1965) No.1, 82–99.

[2] M.Asaoka, D.Turaev, and K.Shinohara, Degenerate behavior in non-hyperbolic semigroup actions on the interval: fast growth of periodic points and universal dynamics, arXiv:1506.07279.

[3] P. Berger, Generic family with robustly infinitely many sinks, preprint, arXiv:1411.6441.

[4] R. Bowen, Topological entropy and axiom A. 1970 Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968) pp. 23–41, Amer. Math. Soc., Providence, R.I.

[5] W. de Melo, A finiteness problem for one dimensional maps. Proe. A.M.S. 101 (1987), 721–727.

[6] Y. Ilyashenko and W. Li, Nonlocal bifurcations. Mathematical Surveys and Monographs, 66. American Mathematical Society, Providence, RI, 1999. xiv+286 pp.

[7] V. Kaloshin, Generic diffeomorphisms with superexponential growth of number of periodic orbits. Comm. Math. Phys. 211 (2000), no. 1, 253–271.

[8] V. Kaloshin and O. S. Kozlovski, A C^r unimodal map with an arbitrary fast growth of the number of periodic points. Ergodic Theory Dynam. Systems 32 (2012), no. 1, 159–165.

[9] V. Kaloshin and M. Saprykina, Generic 3-dimensional volume-preserving diffeomorphisms with superexponential growth of number of periodic orbits. Discrete Contin. Dyn. Syst. 15 (2006), no. 2, 611–640.

[10] O. S. Kozlovski, The dynamics of intersections of analytic manifolds. (Russian) Dokl. Akad. Nauk 323 (1992), no. 5, 823–825; translation in Russian Acad. Sci. Dokl. Math. 45 (1992), no. 2, 425–427 (1993).

[11] S. G. Krantz and H. R. Park, A primer of real analytic functions. Second edition. Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser Boston, Inc., Boston, MA, 2002. xiv+205 pp.

[12] M. Martens, W. de Melo, and S. van Strien, Julia-Fatou-Sullivan theory for real one-dimensional dynamics. Acta Math. 168 (1992), no. 3–4, 273–318.

[13] J. K. Moser, Stable and random motions in dynamical systems. Reprint of the 1973 original. Princeton University Press, Princeton, NJ, 2001. xii+198 pp.

[14] C. L. Siegel and J. K. Moser, Lectures on Celestial Mechanics. Die Grundlehren der mathematischen Wissenschaften, Band 187. Springer-Verlag, New York-Heidelberg (1971).